

## 8.2 Integration by Parts

■ Find an antiderivative using integration by parts.

### Integration by Parts

In this section, you will study an important integration technique called **integration by parts**. This technique can be applied to a wide variety of functions and is particularly useful for integrands involving *products* of algebraic and transcendental functions. For instance, integration by parts works well with integrals such as

$$\int x \ln x \, dx, \quad \int x^2 e^x \, dx, \quad \text{and} \quad \int e^x \sin x \, dx.$$

Integration by parts is based on the formula for the derivative of a product

$$\begin{aligned} \frac{d}{dx}[uv] &= u \frac{dv}{dx} + v \frac{du}{dx} \\ &= uv' + vu' \end{aligned}$$

where both  $u$  and  $v$  are differentiable functions of  $x$ . When  $u'$  and  $v'$  are continuous, you can integrate both sides of this equation to obtain

$$\begin{aligned} uv &= \int uv' \, dx + \int vu' \, dx \\ &= \int u \, dv + \int v \, du. \end{aligned}$$

By rewriting this equation, you obtain the next theorem.

#### THEOREM 8.1 Integration by Parts

If  $u$  and  $v$  are functions of  $x$  and have continuous derivatives, then

$$\int u \, dv = uv - \int v \, du.$$

This formula expresses the original integral in terms of another integral. Depending on the choices of  $u$  and  $dv$ , it may be easier to evaluate the second integral than the original one. Because the choices of  $u$  and  $dv$  are critical in the integration by parts process, the guidelines below are provided.

#### GUIDELINES FOR INTEGRATION BY PARTS


1. Try letting  $dv$  be the most complicated portion of the integrand that fits a basic integration rule. Then  $u$  will be the remaining factor(s) of the integrand.
2. Try letting  $u$  be the portion of the integrand whose derivative is a function simpler than  $u$ . Then  $dv$  will be the remaining factor(s) of the integrand.

Note that  $dv$  always includes the  $dx$  of the original integrand.

When using integration by parts, note that you can first choose  $dv$  or first choose  $u$ . After you choose, however, the choice of the other factor is determined—it must be the remaining portion of the integrand. Also note that  $dv$  must contain the differential  $dx$  of the original integral.

#### Exploration

**Proof Without Words** Here is a different approach to proving the formula for integration by parts. This approach is from “Proof Without Words: Integration by Parts” by Roger B. Nelsen, *Mathematics Magazine*, 64, No. 2, April 1991, p. 130, by permission of the author.



Area   + Area   =  $qs - pr$

$$\int_r^s u \, dv + \int_q^p v \, du = [uv]_{(p,r)}^{(q,s)}$$

$$\int_r^s u \, dv = [uv]_{(p,r)}^{(q,s)} - \int_q^p v \, du$$

Explain how this graph proves the theorem. Which notation in this proof is unfamiliar? What do you think it means?

**EXAMPLE 1** Integration by PartsFind  $\int xe^x dx$ .**Solution** To apply integration by parts, you need to write the integral in the form  $\int u dv$ . There are several ways to do this.

$$\int \underbrace{(x)}_u \underbrace{(e^x dx)}_{dv}, \quad \int \underbrace{(e^x)}_u \underbrace{(x dx)}_{dv}, \quad \int \underbrace{(1)}_u \underbrace{(xe^x dx)}_{dv}, \quad \int \underbrace{(xe^x)}_u \underbrace{(dx)}_{dv}$$

The guidelines on the preceding page suggest the first option because the derivative of  $u = x$  is simpler than  $x$ , and  $dv = e^x dx$  is the most complicated portion of the integrand that fits a basic integration formula.



**REMARK** In Example 1, note that it is not necessary to include a constant of integration when solving

$$v = \int e^x dx = e^x + C_1.$$

To illustrate this, replace  $v = e^x$  by  $v = e^x + C_1$  and apply integration by parts to see that you obtain the same result.

$$dv = e^x dx \Rightarrow v = \int dv = \int e^x dx = e^x$$

$$u = x \Rightarrow du = dx$$

Now, integration by parts produces

$$\begin{aligned} \int u dv &= uv - \int v du && \text{Integration by parts formula} \\ \int xe^x dx &= xe^x - \int e^x dx && \text{Substitute.} \\ &= xe^x - e^x + C. && \text{Integrate.} \end{aligned}$$

To check this, differentiate  $xe^x - e^x + C$  to see that you obtain the original integrand.

**EXAMPLE 2** Integration by PartsFind  $\int x^2 \ln x dx$ .**Solution** In this case,  $x^2$  is more easily integrated than  $\ln x$ . Furthermore, the derivative of  $\ln x$  is simpler than  $\ln x$ . So, you should let  $dv = x^2 dx$ .

$$dv = x^2 dx \Rightarrow v = \int x^2 dx = \frac{x^3}{3}$$

$$u = \ln x \Rightarrow du = \frac{1}{x} dx$$

Integration by parts produces

$$\begin{aligned} \int u dv &= uv - \int v du && \text{Integration by parts formula} \\ \int x^2 \ln x dx &= \frac{x^3}{3} \ln x - \int \left(\frac{x^3}{3}\right)\left(\frac{1}{x}\right) dx && \text{Substitute.} \\ &= \frac{x^3}{3} \ln x - \frac{1}{3} \int x^2 dx && \text{Simplify.} \\ &= \frac{x^3}{3} \ln x - \frac{x^3}{9} + C. && \text{Integrate.} \end{aligned}$$

You can check this result by differentiating.

$$\frac{d}{dx} \left[ \frac{x^3}{3} \ln x - \frac{x^3}{9} + C \right] = \frac{x^3}{3} \left( \frac{1}{x} \right) + (\ln x)(x^2) - \frac{x^2}{3} = x^2 \ln x$$



**TECHNOLOGY** Try graphing

$$\int x^2 \ln x dx \quad \text{and} \quad \frac{x^3}{3} \ln x - \frac{x^3}{9}$$

on your graphing utility. Do you get the same graph? (This may take a while, so be patient.)

One surprising application of integration by parts involves integrands consisting of single terms, such as

$$\int \ln x \, dx \quad \text{or} \quad \int \arcsin x \, dx.$$

In these cases, try letting  $dv = dx$ , as shown in the next example.

### EXAMPLE 3 An Integrand with a Single Term

Evaluate  $\int_0^1 \arcsin x \, dx$ .

**Solution** Let  $dv = dx$ .

$$dv = dx \quad \Rightarrow \quad v = \int dx = x$$

$$u = \arcsin x \quad \Rightarrow \quad du = \frac{1}{\sqrt{1-x^2}} dx$$

Integration by parts produces

$$\int u \, dv = uv - \int v \, du$$

Integration by parts  
formula

$$\begin{aligned} \int \arcsin x \, dx &= x \arcsin x - \int \frac{x}{\sqrt{1-x^2}} dx \\ &= x \arcsin x + \frac{1}{2} \int (1-x^2)^{-1/2} (-2x) dx \\ &= x \arcsin x + \sqrt{1-x^2} + C. \end{aligned}$$


Substitute.

Rewrite.

Integrate.

Using this antiderivative, you can evaluate the definite integral as shown.

$$\begin{aligned} \int_0^1 \arcsin x \, dx &= \left[ x \arcsin x + \sqrt{1-x^2} \right]_0^1 \\ &= \frac{\pi}{2} - 1 \\ &\approx 0.571 \end{aligned}$$



The area of the region is approximately 0.571.

Figure 8.2

The area represented by this definite integral is shown in Figure 8.2.

► **TECHNOLOGY** Remember that there are two ways to use technology to evaluate a definite integral: (1) you can use a numerical approximation such as the Trapezoidal Rule or Simpson's Rule, or (2) you can use a computer algebra system to find the antiderivative and then apply the Fundamental Theorem of Calculus. Both methods have shortcomings. To find the possible error when using a numerical method, the integrand must have a second derivative (Trapezoidal Rule) or a fourth derivative (Simpson's Rule) in the interval of integration; the integrand in Example 3 fails to meet either of these requirements. To apply the Fundamental Theorem of Calculus, the symbolic integration utility must be able to find the antiderivative.

■ **FOR FURTHER INFORMATION** To see how integration by parts is used to prove Stirling's approximation

$$\ln(n!) = n \ln n - n$$

see the article "The Validity of Stirling's Approximation: A Physical Chemistry Project" by A. S. Wallner and K. A. Brandt in *Journal of Chemical Education*.

Some integrals require repeated use of the integration by parts formula.

### EXAMPLE 4 Repeated Use of Integration by Parts

Find  $\int x^2 \sin x \, dx$ .

**Solution** The factors  $x^2$  and  $\sin x$  are equally easy to integrate. However, the derivative of  $x^2$  becomes simpler, whereas the derivative of  $\sin x$  does not. So, you should let  $u = x^2$ .

$$\begin{aligned} dv &= \sin x \, dx &\Rightarrow & v = \int \sin x \, dx = -\cos x \\ u &= x^2 &\Rightarrow & du = 2x \, dx \end{aligned}$$

Now, integration by parts produces

$$\int x^2 \sin x \, dx = -x^2 \cos x + \int 2x \cos x \, dx. \quad \text{First use of integration by parts}$$

This first use of integration by parts has succeeded in simplifying the original integral, but the integral on the right still doesn't fit a basic integration rule. To evaluate that integral, you can apply integration by parts again. This time, let  $u = 2x$ .

$$\begin{aligned} dv &= \cos x \, dx &\Rightarrow & v = \int \cos x \, dx = \sin x \\ u &= 2x &\Rightarrow & du = 2 \, dx \end{aligned}$$

Now, integration by parts produces

$$\begin{aligned} \int 2x \cos x \, dx &= 2x \sin x - \int 2 \sin x \, dx && \text{Second use of integration by parts} \\ &= 2x \sin x + 2 \cos x + C. \end{aligned}$$

Combining these two results, you can write

$$\int x^2 \sin x \, dx = -x^2 \cos x + 2x \sin x + 2 \cos x + C. \quad \blacksquare$$

When making repeated applications of integration by parts, you need to be careful not to interchange the substitutions in successive applications. For instance, in Example 4, the first substitution was  $u = x^2$  and  $dv = \sin x \, dx$ . If, in the second application, you had switched the substitution to  $u = \cos x$  and  $dv = 2x$ , you would have obtained

$$\begin{aligned} \int x^2 \sin x \, dx &= -x^2 \cos x + \int 2x \cos x \, dx \\ &= -x^2 \cos x + x^2 \cos x + \int x^2 \sin x \, dx \\ &= \int x^2 \sin x \, dx \end{aligned}$$

thereby undoing the previous integration and returning to the *original* integral. When making repeated applications of integration by parts, you should also watch for the appearance of a *constant multiple* of the original integral. For instance, this occurs when you use integration by parts to evaluate  $\int e^x \cos 2x \, dx$ , and it also occurs in Example 5 on the next page.

The integral in Example 5 is an important one. In Section 8.4 (Example 5), you will see that it is used to find the arc length of a parabolic segment.

**EXAMPLE 5** Integration by Parts

Find  $\int \sec^3 x \, dx$ .

**Solution** The most complicated portion of the integrand that can be easily integrated is  $\sec^2 x$ , so you should let  $dv = \sec^2 x \, dx$  and  $u = \sec x$ .

$$dv = \sec^2 x \, dx \quad \Rightarrow \quad v = \int \sec^2 x \, dx = \tan x$$

$$u = \sec x \quad \Rightarrow \quad du = \sec x \tan x \, dx$$

Integration by parts produces

$$\int u \, dv = uv - \int v \, du \quad \text{Integration by parts formula}$$

$$\int \sec^3 x \, dx = \sec x \tan x - \int \sec x \tan^2 x \, dx \quad \text{Substitute.}$$

$$\int \sec^3 x \, dx = \sec x \tan x - \int \sec x (\sec^2 x - 1) \, dx \quad \text{Trigonometric identity}$$

$$\int \sec^3 x \, dx = \sec x \tan x - \int \sec^3 x \, dx + \int \sec x \, dx \quad \text{Rewrite.}$$

$$2 \int \sec^3 x \, dx = \sec x \tan x + \int \sec x \, dx \quad \text{Collect like integrals.}$$

$$2 \int \sec^3 x \, dx = \sec x \tan x + \ln|\sec x + \tan x| + C \quad \text{Integrate.}$$

$$\int \sec^3 x \, dx = \frac{1}{2} \sec x \tan x + \frac{1}{2} \ln|\sec x + \tan x| + C. \quad \text{Divide by 2.}$$




Figure 8.3

**EXAMPLE 6** Finding a Centroid

A machine part is modeled by the region bounded by the graph of  $y = \sin x$  and the  $x$ -axis,  $0 \leq x \leq \pi/2$ , as shown in Figure 8.3. Find the centroid of this region.

**Solution** Begin by finding the area of the region.

$$A = \int_0^{\pi/2} \sin x \, dx = \left[ -\cos x \right]_0^{\pi/2} = 1$$

Now, you can find the coordinates of the centroid. To evaluate the integral for  $\bar{y}$ , first rewrite the integrand using the trigonometric identity  $\sin^2 x = (1 - \cos 2x)/2$ .

$$\bar{y} = \frac{1}{A} \int_0^{\pi/2} \frac{\sin x}{2} (\sin x) \, dx = \frac{1}{4} \int_0^{\pi/2} (1 - \cos 2x) \, dx = \frac{1}{4} \left[ x - \frac{\sin 2x}{2} \right]_0^{\pi/2} = \frac{\pi}{8}$$

You can evaluate the integral for  $\bar{x}$ ,  $(1/A) \int_0^{\pi/2} x \sin x \, dx$ , with integration by parts. To do this, let  $dv = \sin x \, dx$  and  $u = x$ . This produces  $v = -\cos x$  and  $du = dx$ , and you can write

$$\int x \sin x \, dx = -x \cos x + \int \cos x \, dx = -x \cos x + \sin x + C.$$

Finally, you can determine  $\bar{x}$  to be

$$\bar{x} = \frac{1}{A} \int_0^{\pi/2} x \sin x \, dx = \left[ -x \cos x + \sin x \right]_0^{\pi/2} = 1.$$

So, the centroid of the region is  $(1, \pi/8)$ .

As you gain experience in using integration by parts, your skill in determining  $u$  and  $dv$  will increase. The next summary lists several common integrals with suggestions for the choices of  $u$  and  $dv$ .



**REMARK** You can use the acronym LIATE as a guideline for choosing  $u$  in integration by parts. In order, check the integrand for the following.

Is there a Logarithmic part?

Is there an Inverse trigonometric part?

Is there an Algebraic part?

Is there a Trigonometric part?

Is there an Exponential part?

### SUMMARY: COMMON INTEGRALS USING INTEGRATION BY PARTS

1. For integrals of the form

$$\int x^n e^{ax} dx, \quad \int x^n \sin ax dx, \quad \text{or} \quad \int x^n \cos ax dx$$

let  $u = x^n$  and let  $dv = e^{ax} dx$ ,  $\sin ax dx$ , or  $\cos ax dx$ .

2. For integrals of the form

$$\int x^n \ln x dx, \quad \int x^n \arcsin ax dx, \quad \text{or} \quad \int x^n \arctan ax dx$$

let  $u = \ln x$ ,  $\arcsin ax$ , or  $\arctan ax$  and let  $dv = x^n dx$ .

3. For integrals of the form

$$\int e^{ax} \sin bx dx \quad \text{or} \quad \int e^{ax} \cos bx dx$$

let  $u = \sin bx$  or  $\cos bx$  and let  $dv = e^{ax} dx$ .

In problems involving repeated applications of integration by parts, a tabular method, illustrated in Example 7, can help to organize the work. This method works well for integrals of the form

$$\int x^n \sin ax dx, \quad \int x^n \cos ax dx, \quad \text{and} \quad \int x^n e^{ax} dx.$$

### EXAMPLE 7 Using the Tabular Method

•••▶ See [LarsonCalculus.com](http://LarsonCalculus.com) for an interactive version of this type of example.

Find  $\int x^2 \sin 4x dx$ .

**Solution** Begin as usual by letting  $u = x^2$  and  $dv = v' dx = \sin 4x dx$ . Next, create a table consisting of three columns, as shown.

Alternate Signs	$u$ and Its Derivatives	$v'$ and Its Antiderivatives
+	$x^2$	$\sin 4x$
−	$2x$	$-\frac{1}{4} \cos 4x$
+	$2$	$-\frac{1}{16} \sin 4x$
−	$0$	$\frac{1}{64} \cos 4x$

↑  
Differentiate until you obtain 0 as a derivative.

The solution is obtained by adding the signed products of the diagonal entries:

$$\int x^2 \sin 4x dx = -\frac{1}{4} x^2 \cos 4x + \frac{1}{8} x \sin 4x + \frac{1}{32} \cos 4x + C.$$

### FOR FURTHER INFORMATION

For more information on the tabular method, see the article “Tabular Integration by Parts” by David Horowitz in *The College Mathematics Journal*, and the article “More on Tabular Integration by Parts” by Leonard Gillman in *The College Mathematics Journal*. To view these articles, go to [MathArticles.com](http://MathArticles.com).

## 8.2 Exercises

See [CalcChat.com](http://CalcChat.com) for tutorial help and worked-out solutions to odd-numbered exercises.

**Setting Up Integration by Parts** In Exercises 1–6, identify  $u$  and  $dv$  for finding the integral using integration by parts. (Do not evaluate the integral.)

1.  $\int x e^{2x} dx$
2.  $\int x^2 e^{2x} dx$
3.  $\int (\ln x)^2 dx$
4.  $\int \ln 5x dx$
5.  $\int x \sec^2 x dx$
6.  $\int x^2 \cos x dx$

**Using Integration by Parts** In Exercises 7–10, evaluate the integral using integration by parts with the given choices of  $u$  and  $dv$ .

7.  $\int x^3 \ln x dx$ ;  $u = \ln x$ ,  $dv = x^3 dx$
8.  $\int (4x + 7)e^x dx$ ;  $u = 4x + 7$ ,  $dv = e^x dx$
9.  $\int x \sin 3x dx$ ;  $u = x$ ,  $dv = \sin 3x dx$
10.  $\int x \cos 4x dx$ ;  $u = x$ ,  $dv = \cos 4x dx$

**Finding an Indefinite Integral** In Exercises 11–30, find the indefinite integral. (Note: Solve by the simplest method—not all require integration by parts.)

11.  $\int x e^{-4x} dx$
12.  $\int \frac{5x}{e^{2x}} dx$
13.  $\int x^3 e^x dx$
14.  $\int \frac{e^{1/t}}{t^2} dt$
15.  $\int t \ln(t + 1) dt$
16.  $\int x^5 \ln 3x dx$
17.  $\int \frac{(\ln x)^2}{x} dx$
18.  $\int \frac{\ln x}{x^3} dx$
19.  $\int \frac{x e^{2x}}{(2x + 1)^2} dx$
20.  $\int \frac{x^3 e^{x^2}}{(x^2 + 1)^2} dx$
21.  $\int x \sqrt{x - 5} dx$
22.  $\int \frac{x}{\sqrt{6x + 1}} dx$
23.  $\int x \cos x dx$
24.  $\int t \csc t \cot t dt$
25.  $\int x^3 \sin x dx$
26.  $\int x^2 \cos x dx$
27.  $\int \arctan x dx$
28.  $\int 4 \arccos x dx$
29.  $\int e^{-3x} \sin 5x dx$
30.  $\int e^{4x} \cos 2x dx$

**Differential Equation** In Exercises 31–34, solve the differential equation.

31.  $y' = \ln x$
32.  $y' = \arctan \frac{x}{2}$



$$33. \frac{dy}{dt} = \frac{t^2}{\sqrt{3 + 5t}}$$

$$34. \frac{dy}{dx} = x^2 \sqrt{x - 3}$$



**Slope Field** In Exercises 35 and 36, a differential equation, a point, and a slope field are given. (a) Sketch two approximate solutions of the differential equation on the slope field, one of which passes through the given point. (b) Use integration to find the particular solution of the differential equation and use a graphing utility to graph the solution. Compare the result with the sketches in part (a). To print an enlarged copy of the graph, go to [MathGraphs.com](http://MathGraphs.com).

$$35. \frac{dy}{dx} = x\sqrt{y} \cos x, (0, 4) \quad 36. \frac{dy}{dx} = e^{-x/3} \sin 2x, (0, -\frac{18}{37})$$



**Slope Field** In Exercises 37 and 38, use a computer algebra system to graph the slope field for the differential equation and graph the solution through the specified initial condition.

$$37. \frac{dy}{dx} = \frac{x}{y} e^{x/8}, y(0) = 2 \quad 38. \frac{dy}{dx} = \frac{x}{y} \sin x, y(0) = 4$$

**Evaluating a Definite Integral** In Exercises 39–48, evaluate the definite integral. Use a graphing utility to confirm your result.

39.  $\int_0^3 x e^{x/2} dx$
40.  $\int_0^2 x^2 e^{-2x} dx$
41.  $\int_0^{\pi/4} x \cos 2x dx$
42.  $\int_0^{\pi} x \sin 2x dx$
43.  $\int_0^{1/2} \arccos x dx$
44.  $\int_0^1 x \arcsin x^2 dx$
45.  $\int_0^1 e^x \sin x dx$
46.  $\int_0^1 \ln(4 + x^2) dx$
47.  $\int_2^4 x \operatorname{arcsec} x dx$
48.  $\int_0^{\pi/8} x \sec^2 2x dx$

**Using the Tabular Method** In Exercises 49–54, use the tabular method to find the integral.

49.  $\int x^2 e^{2x} dx$
50.  $\int x^3 e^{-2x} dx$
51.  $\int x^3 \sin x dx$
52.  $\int x^3 \cos 2x dx$

53.  $\int x \sec^2 x \, dx$

54.  $\int x^2(x-2)^{3/2} \, dx$

**Using Two Methods Together** In Exercises 55–58, find the indefinite integral by using substitution followed by integration by parts.

55.  $\int \sin \sqrt{x} \, dx$

56.  $\int 2x^3 \cos x^2 \, dx$

57.  $\int x^5 e^{x^2} \, dx$

58.  $\int e^{\sqrt{2x}} \, dx$

### WRITING ABOUT CONCEPTS

#### 59. Integration by Parts

- (a) Integration by parts is based on what differentiation rule? Explain.  
 (b) In your own words, state how you determine which parts of the integrand should be  $u$  and  $dv$ .

**60. Integration by Parts** When evaluating  $\int x \sin x \, dx$ , explain how letting  $u = \sin x$  and  $dv = x \, dx$  makes the solution more difficult to find.

**61. Integration by Parts** State whether you would use integration by parts to evaluate each integral. If so, identify what you would use for  $u$  and  $dv$ . Explain your reasoning.

- (a)  $\int \frac{\ln x}{x} \, dx$  (b)  $\int x \ln x \, dx$  (c)  $\int x^2 e^{-3x} \, dx$   
 (d)  $\int 2xe^{x^2} \, dx$  (e)  $\int \frac{x}{\sqrt{x+1}} \, dx$  (f)  $\int \frac{x}{\sqrt{x^2+1}} \, dx$

**64. Using Two Methods** Integrate  $\int x\sqrt{4-x} \, dx$

- (a) by parts, letting  $dv = \sqrt{4-x} \, dx$ .  
 (b) by substitution, letting  $u = 4-x$ .



**Finding a General Rule** In Exercises 65 and 66, use a computer algebra system to find the integrals for  $n = 0, 1, 2$ , and 3. Use the result to obtain a general rule for the integrals for any positive integer  $n$  and test your results for  $n = 4$ .

65.  $\int x^n \ln x \, dx$

66.  $\int x^n e^x \, dx$

**Proof** In Exercises 67–72, use integration by parts to prove the formula. (For Exercises 67–70, assume that  $n$  is a positive integer.)

67.  $\int x^n \sin x \, dx = -x^n \cos x + n \int x^{n-1} \cos x \, dx$

68.  $\int x^n \cos x \, dx = x^n \sin x - n \int x^{n-1} \sin x \, dx$

69.  $\int x^n \ln x \, dx = \frac{x^{n+1}}{(n+1)^2} [-1 + (n+1) \ln x] + C$

70.  $\int x^n e^{ax} \, dx = \frac{x^n e^{ax}}{a} - \frac{n}{a} \int x^{n-1} e^{ax} \, dx$

71.  $\int e^{ax} \sin bx \, dx = \frac{e^{ax}(a \sin bx - b \cos bx)}{a^2 + b^2} + C$

72.  $\int e^{ax} \cos bx \, dx = \frac{e^{ax}(a \cos bx + b \sin bx)}{a^2 + b^2} + C$

**Using Formulas** In Exercises 73–78, find the integral by using the appropriate formula from Exercises 67–72.

73.  $\int x^2 \sin x \, dx$

74.  $\int x^2 \cos x \, dx$

75.  $\int x^5 \ln x \, dx$

76.  $\int x^3 e^{2x} \, dx$

77.  $\int e^{-3x} \sin 4x \, dx$

78.  $\int e^{2x} \cos 3x \, dx$



**Area** In Exercises 79–82, use a graphing utility to graph the region bounded by the graphs of the equations. Then find the area of the region analytically.

79.  $y = 2xe^{-x}, y = 0, x = 3$

80.  $y = \frac{1}{10}xe^{3x}, y = 0, x = 0, x = 2$

81.  $y = e^{-x} \sin \pi x, y = 0, x = 1$


82.  $y = x^3 \ln x, y = 0, x = 1, x = 3$

**83. Area, Volume, and Centroid** Given the region bounded by the graphs of  $y = \ln x, y = 0$ , and  $x = e$ , find

- (a) the area of the region.  
 (b) the volume of the solid generated by revolving the region about the  $x$ -axis.  
 (c) the volume of the solid generated by revolving the region about the  $y$ -axis.  
 (d) the centroid of the region.



**62. HOW DO YOU SEE IT?** Use the graph of  $f'$  shown in the figure to answer the following.



- (a) Approximate the slope of  $f$  at  $x = 2$ . Explain.  
 (b) Approximate any open intervals in which the graph of  $f$  is increasing and any open intervals in which it is decreasing. Explain.

**63. Using Two Methods** Integrate  $\int \frac{x^3}{\sqrt{4+x^2}} \, dx$

- (a) by parts, letting  $dv = \frac{x}{\sqrt{4+x^2}} \, dx$ .  
 (b) by substitution, letting  $u = 4 + x^2$ .



- 84. Area, Volume, and Centroid** Given the region bounded by the graphs of  $y = x \sin x$ ,  $y = 0$ ,  $x = 0$ , and  $x = \pi$ , find
- the area of the region.
  - the volume of the solid generated by revolving the region about the  $x$ -axis.
  - the volume of the solid generated by revolving the region about the  $y$ -axis.
  - the centroid of the region.

**85. Centroid** Find the centroid of the region bounded by the graphs of  $y = \arcsin x$ ,  $x = 0$ , and  $y = \pi/2$ . How is this problem related to Example 6 in this section?

**86. Centroid** Find the centroid of the region bounded by the graphs of  $f(x) = x^2$ ,  $g(x) = 2^x$ ,  $x = 2$ , and  $x = 4$ .

**87. Average Displacement** A damping force affects the vibration of a spring so that the displacement of the spring is given by

$$y = e^{-4t} (\cos 2t + 5 \sin 2t).$$

Find the average value of  $y$  on the interval from  $t = 0$  to  $t = \pi$ .

• • **88. Memory Model** • • • • •

A model for the ability  $M$  of a child to memorize, measured on a scale from 0 to 10, is given by

$$M = 1 + 1.6t \ln t, \quad 0 < t \leq 4$$

where  $t$  is the child's age in years. Find the average value of this model

- between the child's first and second birthdays.
- between the child's third and fourth birthdays.



**Present Value** In Exercises 89 and 90, find the present value  $P$  of a continuous income flow of  $c(t)$  dollars per year for

$$P = \int_0^{t_1} c(t)e^{-rt} dt$$

where  $t_1$  is the time in years and  $r$  is the annual interest rate compounded continuously.

**89.**  $c(t) = 100,000 + 4000t$ ,  $r = 5\%$ ,  $t_1 = 10$

**90.**  $c(t) = 30,000 + 500t$ ,  $r = 7\%$ ,  $t_1 = 5$

**Integrals Used to Find Fourier Coefficients** In Exercises 91 and 92, verify the value of the definite integral, where  $n$  is a positive integer.

$$91. \int_{-\pi}^{\pi} x \sin nx \, dx = \begin{cases} \frac{2\pi}{n}, & n \text{ is odd} \\ -\frac{2\pi}{n}, & n \text{ is even} \end{cases}$$

$$92. \int_{-\pi}^{\pi} x^2 \cos nx \, dx = \frac{(-1)^n 4\pi}{n^2}$$

**93. Vibrating String** A string stretched between the two points  $(0, 0)$  and  $(2, 0)$  is plucked by displacing the string  $h$  units at its midpoint. The motion of the string is modeled by a **Fourier Sine Series** whose coefficients are given by

$$b_n = h \int_0^1 x \sin \frac{n\pi x}{2} dx + h \int_1^2 (-x + 2) \sin \frac{n\pi x}{2} dx.$$

Find  $b_n$ .



**94. Euler's Method** Consider the differential equation  $f'(x) = xe^{-x}$  with the initial condition  $f(0) = 0$ .

- Use integration to solve the differential equation.
- Use a graphing utility to graph the solution of the differential equation.
- Use Euler's Method with  $h = 0.05$ , and the recursive capabilities of a graphing utility, to generate the first 80 points of the graph of the approximate solution. Use the graphing utility to plot the points. Compare the result with the graph in part (b).
- Repeat part (c) using  $h = 0.1$  and generate the first 40 points.
- Why is the result in part (c) a better approximation of the solution than the result in part (d)?



**Euler's Method** In Exercises 95 and 96, consider the differential equation and repeat parts (a)–(d) of Exercise 94.

**95.**  $f'(x) = 3x \sin(2x)$   
 $f(0) = 0$

**96.**  $f'(x) = \cos \sqrt{x}$   
 $f(0) = 1$

**97. Think About It** Give a geometric explanation of why

$$\int_0^{\pi/2} x \sin x \, dx \leq \int_0^{\pi/2} x \, dx.$$

Verify the inequality by evaluating the integrals.

**98. Finding a Pattern** Find the area bounded by the graphs of  $y = x \sin x$  and  $y = 0$  over each interval.

(a)  $[0, \pi]$  (b)  $[\pi, 2\pi]$  (c)  $[2\pi, 3\pi]$

Describe any patterns that you notice. What is the area between the graphs of  $y = x \sin x$  and  $y = 0$  over the interval  $[n\pi, (n+1)\pi]$ , where  $n$  is any nonnegative integer? Explain.

**99. Finding an Error** Find the fallacy in the following argument that  $0 = 1$ .

$$dv = dx \Rightarrow v = \int dx = x$$

$$u = \frac{1}{x} \Rightarrow du = -\frac{1}{x^2} dx$$

$$0 + \int \frac{dx}{x} = \left(\frac{1}{x}\right)(x) - \int \left(-\frac{1}{x^2}\right)(x) dx \\ = 1 + \int \frac{dx}{x}$$

$$\text{So, } 0 = 1.$$

Juriah Mosin/Shutterstock.com